

1 Evaluate

a $\sqrt{49}$ b $\sqrt{121}$ c $\sqrt{\frac{1}{9}}$ d $\sqrt{\frac{4}{25}}$ e $\sqrt{0.01}$ f $\sqrt{0.09}$
 g $\sqrt[3]{8}$ h $\sqrt[3]{1000}$ i $\sqrt[4]{81}$ j $\sqrt[4]{1\frac{9}{16}}$ k $\sqrt[3]{0.125}$ l $\sqrt[3]{15\frac{5}{8}}$

2 Simplify

a $\sqrt{7} \times \sqrt{7}$ b $4\sqrt{5} \times \sqrt{5}$ c $(3\sqrt{3})^2$ d $(\sqrt{6})^4$
 e $(\sqrt{2})^5$ f $(2\sqrt{3})^3$ g $\sqrt{2} \times \sqrt{8}$ h $2\sqrt{3} \times \sqrt{27}$
 i $\frac{\sqrt{32}}{\sqrt{2}}$ j $\frac{\sqrt{3}}{\sqrt{12}}$ k $(\sqrt[3]{6})^3$ l $(3\sqrt[3]{2})^3$

3 Express in the form $k\sqrt{2}$

a $\sqrt{18}$ b $\sqrt{50}$ c $\sqrt{8}$ d $\sqrt{98}$ e $\sqrt{200}$ f $\sqrt{162}$

4 Simplify

a $\sqrt{12}$ b $\sqrt{28}$ c $\sqrt{80}$ d $\sqrt{27}$ e $\sqrt{24}$ f $\sqrt{128}$
 g $\sqrt{45}$ h $\sqrt{40}$ i $\sqrt{75}$ j $\sqrt{112}$ k $\sqrt{99}$ l $\sqrt{147}$
 m $\sqrt{216}$ n $\sqrt{800}$ o $\sqrt{180}$ p $\sqrt{60}$ q $\sqrt{363}$ r $\sqrt{208}$

5 Simplify

a $\sqrt{18} + \sqrt{50}$ b $\sqrt{48} - \sqrt{27}$ c $2\sqrt{8} + \sqrt{72}$
 d $\sqrt{360} - 2\sqrt{40}$ e $2\sqrt{5} - \sqrt{45} + 3\sqrt{20}$ f $\sqrt{24} + \sqrt{150} - 2\sqrt{96}$

6 Express in the form $a + b\sqrt{3}$

a $\sqrt{3}(2 + \sqrt{3})$ b $4 - \sqrt{3} - 2(1 - \sqrt{3})$ c $(1 + \sqrt{3})(2 + \sqrt{3})$
 d $(4 + \sqrt{3})(1 + 2\sqrt{3})$ e $(3\sqrt{3} - 4)^2$ f $(3\sqrt{3} + 1)(2 - 5\sqrt{3})$

7 Simplify

a $(\sqrt{5} + 1)(2\sqrt{5} + 3)$ b $(1 - \sqrt{2})(4\sqrt{2} - 3)$ c $(2\sqrt{7} + 3)^2$
 d $(3\sqrt{2} - 1)(2\sqrt{2} + 5)$ e $(\sqrt{5} - \sqrt{2})(\sqrt{5} + 2\sqrt{2})$ f $(3 - \sqrt{8})(4 + \sqrt{2})$

8 Express each of the following as simply as possible with a rational denominator.

a $\frac{1}{\sqrt{5}}$ b $\frac{2}{\sqrt{3}}$ c $\frac{1}{\sqrt{8}}$ d $\frac{14}{\sqrt{7}}$ e $\frac{3\sqrt{2}}{\sqrt{3}}$ f $\frac{\sqrt{5}}{\sqrt{15}}$
 g $\frac{1}{3\sqrt{7}}$ h $\frac{12}{\sqrt{72}}$ i $\frac{1}{\sqrt{80}}$ j $\frac{3}{2\sqrt{54}}$ k $\frac{4\sqrt{20}}{3\sqrt{18}}$ l $\frac{3\sqrt{175}}{2\sqrt{27}}$

9 Simplify

a $\sqrt{8} + \frac{6}{\sqrt{2}}$

b $\sqrt{48} - \frac{10}{\sqrt{3}}$

c $\frac{6-\sqrt{8}}{\sqrt{2}}$

d $\frac{\sqrt{45}-5}{\sqrt{20}}$

e $\frac{1}{\sqrt{18}} + \frac{1}{\sqrt{32}}$

f $\frac{2}{\sqrt{3}} - \frac{\sqrt{6}}{\sqrt{72}}$

10 Solve each equation, giving your answers as simply as possible in terms of surds.

a $x(x+4) = 4(x+8)$

b $x - \sqrt{48} = 2\sqrt{3} - 2x$

c $x\sqrt{18} - 4 = \sqrt{8}$

d $x\sqrt{5} + 2 = \sqrt{20}(x-1)$

11 a Simplify $(2 - \sqrt{3})(2 + \sqrt{3})$.

b Express $\frac{2}{2-\sqrt{3}}$ in the form $a + b\sqrt{3}$.

12 Express each of the following as simply as possible with a rational denominator.

a $\frac{1}{\sqrt{2}+1}$

b $\frac{4}{\sqrt{3}-1}$

c $\frac{1}{\sqrt{6}-2}$

d $\frac{3}{2+\sqrt{3}}$

e $\frac{1}{2+\sqrt{5}}$

f $\frac{\sqrt{2}}{\sqrt{2}-1}$

g $\frac{6}{\sqrt{7}+3}$

h $\frac{1}{3+2\sqrt{2}}$

i $\frac{1}{4-2\sqrt{3}}$

j $\frac{3}{3\sqrt{2}+4}$

k $\frac{2\sqrt{3}}{7-4\sqrt{3}}$


l $\frac{6}{\sqrt{5}-\sqrt{3}}$

13 Solve the equation

$$3x = \sqrt{5}(x+2),$$

giving your answer in the form $a + b\sqrt{5}$, where a and b are rational.

14



The diagram shows a rectangle measuring $(3\sqrt{2} - 3) \text{ cm}$ by $l \text{ cm}$.

Given that the area of the rectangle is 6 cm^2 , find the exact value of l in its simplest form.

15 Express each of the following as simply as possible with a rational denominator.

a $\frac{\sqrt{2}}{\sqrt{2}+\sqrt{6}}$

b $\frac{1+\sqrt{3}}{2+\sqrt{3}}$

c $\frac{1+\sqrt{10}}{\sqrt{10}-3}$

d $\frac{3-\sqrt{2}}{4+3\sqrt{2}}$

e $\frac{1-\sqrt{2}}{3-\sqrt{8}}$

f $\frac{\sqrt{3}-5}{2\sqrt{3}-4}$

g $\frac{\sqrt{12}+3}{3-\sqrt{3}}$

h $\frac{3\sqrt{7}-2}{2\sqrt{7}-5}$

1 Evaluate

a 8^2 **b** 6^3 **c** 7^0 **d** $(-5)^4$ **e** $(-3)^5$ **f** $(\frac{1}{2})^4$
g $(\frac{2}{3})^3$ **h** $(-\frac{1}{4})^3$ **i** $(1\frac{1}{3})^2$ **j** $(1\frac{1}{2})^4$ **k** $(0.1)^5$ **l** $(-0.2)^3$

2 Write in the form 2^n

a $2^5 \times 2^3$ **b** 2×2^6 **c** 1 **d** $2^6 \div 2^2$ **e** $2^{15} \div 2^6$ **f** $(2^7)^2$

3 Simplify

a $2p^2 \times 4p^5$ **b** $x^2 \times x^3 \times x^5$ **c** $12n^7 \div 2n^2$ **d** $(y^3)^4$
e $(2b)^3 \div 4b^2$ **f** $p^3q \times pq^2$ **g** $x^4y^3 \div xy^2$ **h** $2r^2s \times 3s^2$
i $6x^5y^8 \div 3x^2y$ **j** $6a^4b^5 \times \frac{2}{3}ab^3$ **k** $(5rs^2)^3 \div (10rs)^2$ **l** $3p^4q^3 \div \frac{1}{5}pq^2$

4 Evaluate

a 3^{-2} **b** $(\frac{2}{5})^0$ **c** $(-2)^{-6}$ **d** $(\frac{1}{6})^{-2}$ **e** $(1\frac{1}{2})^{-3}$ **f** $9^{\frac{1}{2}}$
g $16^{\frac{1}{4}}$ **h** $(-27)^{\frac{1}{3}}$ **i** $(\frac{1}{49})^{\frac{1}{2}}$ **j** $125^{\frac{1}{3}}$ **k** $(\frac{4}{9})^{\frac{1}{2}}$ **l** $36^{-\frac{1}{2}}$
m $81^{-\frac{1}{4}}$ **n** $(-64)^{-\frac{1}{3}}$ **o** $(\frac{1}{32})^{-\frac{1}{5}}$ **p** $(-\frac{8}{125})^{\frac{1}{3}}$ **q** $(2\frac{1}{4})^{\frac{1}{2}}$ **r** $(3\frac{3}{8})^{-\frac{1}{3}}$

5 Evaluate

a $4^{\frac{3}{2}}$ **b** $27^{\frac{2}{3}}$ **c** $16^{\frac{3}{4}}$ **d** $(-125)^{\frac{2}{3}}$ **e** $9^{\frac{5}{2}}$ **f** $8^{-\frac{2}{3}}$
g $36^{-\frac{3}{2}}$ **h** $(\frac{1}{8})^{\frac{4}{3}}$ **i** $(\frac{4}{9})^{\frac{3}{2}}$ **j** $(\frac{1}{216})^{-\frac{2}{3}}$ **k** $(\frac{9}{16})^{-\frac{3}{2}}$ **l** $(-\frac{27}{64})^{\frac{4}{3}}$
m $(0.04)^{\frac{1}{2}}$ **n** $(2.25)^{-\frac{3}{2}}$ **o** $(0.064)^{\frac{2}{3}}$ **p** $(1\frac{9}{16})^{-\frac{3}{2}}$ **q** $(5\frac{1}{16})^{\frac{3}{4}}$ **r** $(2\frac{10}{27})^{-\frac{4}{3}}$

6 Work out

a $4^{\frac{1}{2}} \times 27^{\frac{1}{3}}$ **b** $16^{\frac{1}{4}} + 25^{\frac{1}{2}}$ **c** $8^{-\frac{1}{3}} \div 36^{\frac{1}{2}}$ **d** $(-64)^{\frac{1}{3}} \times 9^{\frac{3}{2}}$
e $(\frac{1}{3})^{-2} - (-8)^{\frac{1}{3}}$ **f** $(\frac{1}{25})^{\frac{1}{2}} \times (\frac{1}{4})^{-2}$ **g** $81^{\frac{3}{4}} - (\frac{1}{49})^{-\frac{1}{2}}$ **h** $(\frac{1}{27})^{-\frac{1}{3}} \times (\frac{4}{9})^{-\frac{3}{2}}$
i $(\frac{1}{9})^{-\frac{1}{2}} \times (-32)^{\frac{3}{5}}$ **j** $(121)^{0.5} + (32)^{0.2}$ **k** $(100)^{0.5} \div (0.25)^{1.5}$ **l** $(16)^{-0.25} \times (243)^{0.4}$

7 Simplify

a $x^8 \times x^{-6}$ **b** $y^{-2} \times y^{-4}$ **c** $6p^3 \div 2p^7$ **d** $(2x^{-4})^3$
e $y^3 \times y^{-\frac{1}{2}}$ **f** $2b^{\frac{2}{3}} \times 4b^{\frac{1}{4}}$ **g** $x^{\frac{3}{5}} \div x^{\frac{1}{3}}$ **h** $a^{\frac{1}{2}} \div a^{\frac{4}{3}}$
i $p^{\frac{1}{4}} \div p^{-\frac{1}{5}}$ **j** $(3x^{\frac{2}{5}})^2$ **k** $y \times y^{\frac{5}{6}} \times y^{-\frac{3}{2}}$ **l** $4t^{\frac{3}{2}} \div 12t^{\frac{1}{2}}$
m $\frac{b^2 \times b^{\frac{1}{4}}}{b^{\frac{1}{2}}}$ **n** $\frac{y^{\frac{1}{2}} \times y^{\frac{1}{3}}}{y}$ **o** $\frac{4x^{\frac{2}{3}} \times 3x^{-\frac{1}{6}}}{6x^{\frac{3}{4}}}$ **p** $\frac{2a \times a^{\frac{3}{4}}}{8a^{-\frac{1}{2}}}$

8 Solve each equation.

a $x^{\frac{1}{2}} = 6$

b $x^{\frac{1}{3}} = 5$

c $x^{-\frac{1}{2}} = 2$

d $x^{-\frac{1}{4}} = \frac{1}{3}$

e $x^{\frac{3}{2}} = 8$

f $x^{\frac{2}{3}} = 16$

g $x^{\frac{4}{3}} = 81$

h $x^{-\frac{3}{2}} = 27$

9 Express in the form x^k

a \sqrt{x}

b $\frac{1}{\sqrt[3]{x}}$

c $x^2 \times \sqrt{x}$

d $\frac{\sqrt[4]{x}}{x}$

e $\sqrt{x^3}$

f $\sqrt{x} \times \sqrt[3]{x}$

g $(\sqrt{x})^5$

h $\sqrt[3]{x^2} \times (\sqrt{x})^3$

10 Express each of the following in the form ax^b , where a and b are rational constants.

a $\frac{4}{\sqrt{x}}$

b $\frac{1}{2x}$

c $\frac{3}{4x^3}$

d $\frac{1}{(3x)^2}$

e $\frac{2}{5\sqrt[3]{x}}$

f $\frac{1}{\sqrt{9x^3}}$

11 Express in the form 2^k

a 8^2

b $(\frac{1}{4})^{-2}$

c $(\frac{1}{2})^{\frac{1}{3}}$

d $16^{-\frac{1}{6}}$

e $8^{\frac{2}{5}}$

f $(\frac{1}{32})^{-3}$

12 Express each of the following in the form 3^y , where y is a function of x .

a 9^x

b 81^{x+1}

c $27^{\frac{x}{3}}$

d $(\frac{1}{3})^x$

e 9^{2x-1}

f $(\frac{1}{27})^{x+2}$

13 Given that $y = 2^x$, express each of the following in terms of y .

a 2^{x+1}

b 2^{x-2}

c 2^{2x}

d 8^x

e 2^{4x+3}

f $(\frac{1}{2})^{x-3}$

14 Find the value of x such that

a $2^x = 64$

b $5^{x-1} = 125$

c $3^{x+4} - 27 = 0$

d $8^x - 2 = 0$

e $3^{2x-1} = 9$

f $16 - 4^{3x-2} = 0$

g $9^{x-2} = 27$

h $8^{2x+1} = 16$

i $49^{x+1} = \sqrt{7}$

j $3^{3x-2} = \sqrt[3]{9}$

k $(\frac{1}{6})^{x+3} = 36$

l $(\frac{1}{2})^{3x-1} = 8$

15 Solve each equation.

a $2^{x+3} = 4^x$

b $5^{3x} = 25^{x+1}$

c $9^{2x} = 3^{x-3}$

d $16^x = 4^{1-x}$

e $4^{x+2} = 8^x$

f $27^{2x} = 9^{3-x}$

g $6^{3x-1} = 36^{x+2}$

h $8^x = 16^{2x-1}$

i $125^x = 5^{x-3}$

j $(\frac{1}{3})^x = 3^{x-4}$

k $(\frac{1}{2})^{1-x} = (\frac{1}{8})^{2x}$

l $(\frac{1}{4})^{x+1} = 8^x$

16 Expand and simplify

a $x(x^2 - x^{-1})$

b $2x^3(x^{-1} + 3)$

c $x^{-1}(3x - x^3)$

d $4x^{-2}(3x^5 + 2x^3)$

e $\frac{1}{2}x^2(6x + 4x^{-1})$

f $3x^{\frac{1}{2}}(x^{-\frac{1}{2}} - x^{\frac{3}{2}})$

g $x^{-\frac{3}{2}}(5x^2 + x^{\frac{7}{2}})$

h $x^{\frac{1}{3}}(3x^{\frac{5}{3}} - x^{-\frac{4}{3}})$

i $(x^2 + 1)(x^4 - 3)$

j $(2x^5 + x)(x^4 + 3)$

k $(x^2 - 2x^{-1})(x - x^{-2})$

l $(x^2 - x^{\frac{3}{2}})(x - x^{\frac{1}{2}})$

17 Simplify

a $\frac{x^3 + 2x}{x}$

b $\frac{4t^5 - 6t^3}{2t^2}$

c $\frac{x^{\frac{3}{2}} - 3x}{x^{\frac{1}{2}}}$

d $\frac{y^2(y^3 - 6)}{3y}$

e $\frac{p + p^{\frac{3}{2}}}{p^{\frac{3}{4}}}$

f $\frac{8w - 2w^{\frac{1}{2}}}{4w^{-\frac{1}{2}}}$

g $\frac{x+1}{x^{\frac{1}{2}} + x^{-\frac{1}{2}}}$

h $\frac{2t^3 - 4t}{t^{\frac{3}{2}} - 2t^{-\frac{1}{2}}}$

- 1 Express each of the following in the form $a\sqrt{2} + b\sqrt{3}$, where a and b are integers.

a $\sqrt{27} + 2\sqrt{50}$

b $\sqrt{6}(\sqrt{3} - \sqrt{8})$

- 2 Given that $x > 0$, find in the form $k\sqrt{3}$ the value of x such that

$$x(x - 2) = 2(6 - x).$$

- 3 Solve the equation

$$25^x = 5^{4x+1}.$$

- 4 a Express $\sqrt[3]{24}$ in the form $k\sqrt[3]{3}$.

- b Find the integer n such that

$$\sqrt[3]{24} + \sqrt[3]{81} = \sqrt[3]{n}.$$

- 5 Show that

$$\frac{10\sqrt{3}}{\sqrt{15}} + \frac{4}{\sqrt{5}-\sqrt{7}}$$

can be written in the form $k\sqrt{7}$, where k is an integer to be found.

- 6 Showing your method clearly,

a express $\sqrt{37.5}$ in the form $a\sqrt{6}$,

b express $\sqrt{9\frac{3}{5}} - \sqrt{6\frac{2}{3}}$ in the form $b\sqrt{15}$.

- 7 Given that $x = 2^{t-1}$ and $y = 2^{3t}$,

- a find expressions in terms of t for

i xy ii $2y^2$

- b Hence, or otherwise, find the value of t for which

$$2y^2 - xy = 0.$$

- 8 Solve the equation

$$\sqrt{2}(3x - 1) = 2(2x + 3),$$

giving your answer in the form $a + b\sqrt{2}$, where a and b are integers.

- 9 Given that $6^{y+1} = 36^{x-2}$,

- a express y in the form $ax + b$,

- b find the value of $4^{x - \frac{1}{2}y}$.

- 10 Express each of the following in the form $a + b\sqrt{2}$, where a and b are integers.

a $(3 - \sqrt{2})(1 + \sqrt{2})$

b $\frac{\sqrt{2}}{\sqrt{2}-1}$

- 11 Solve the equation

$$16^{x+1} = 8^{2x+1}.$$

- 12 Given that

$$(a - 2\sqrt{3})^2 = b - 20\sqrt{3},$$

find the values of the integers a and b .

- 13 a Find the value of
- t
- such that

$$\left(\frac{1}{4}\right)^{t-3} = 8.$$

- b Solve the equation

$$\left(\frac{1}{3}\right)^y = 27^{y+1}.$$

- 14 Express each of the following in the form
- $a + b\sqrt{5}$
- , where
- a
- and
- b
- are integers.

a $\sqrt{20}(\sqrt{5} - 3)$

b $(1 - \sqrt{5})(3 + 2\sqrt{5})$


c $\frac{1+\sqrt{5}}{\sqrt{5}-2}$

- 15 Given that
- $a^{\frac{1}{3}} = b^{\frac{3}{4}}$
- , and that
- $a > 0$
- and
- $b > 0$
- ,

- a find an expression for
- $a^{\frac{1}{2}}$
- in terms of
- b
- ,

- b find an expression for
- $b^{\frac{1}{2}}$
- in terms of
- a
- .

- 16



In triangle ABC , $AB = 2\sqrt{3} - 1$, $BC = \sqrt{3} + 2$ and $\angle ABC = 90^\circ$.

- a Find the exact area of triangle
- ABC
- in its simplest form.

- b Show that
- $AC = 2\sqrt{5}$
- .

- c Show that
- $\tan(\angle ACB) = 5\sqrt{3} - 8$
- .

- 17 a Given that
- $y = 2^x$
- , express each of the following in terms of
- y
- .

i 2^{x+2}

ii 4^x

- b Hence, or otherwise, find the value of
- x
- for which

$$4^x - 2^{x+2} = 0.$$

- 18 Given that the point with coordinates
- $(1 + \sqrt{3}, 5\sqrt{3})$
- lies on the curve with the equation

$$y = 2x^2 + px + q,$$

find the values of the rational constants p and q .

1 Giving your answers in descending powers of x , simplify

a $(x^2 + 3x + 2) + (2x^2 + 5x + 1)$

b $(x^3 + 4x^2 + x - 6) + (x^2 - 3x + 7)$

c $(4 - x + 2x^3) + (3 - x + 6x^2 - 5x^3)$

d $(x^5 + 8x^3 - 5x^2 - 9) + (-x^4 - 4x + 1)$

e $(3x^3 - 7x^2 + 2) - (x^3 + 2x^2 + x - 6)$

f $(x^5 + 3x^4 - x^2 - 3) - (x^4 + 2x^3 - 3x + 2)$

g $(2x^7 - 9x^5 + x^3 + x) - (3x^6 - 4x^3 + x + 5)$

h $2(x^4 + 4x^2 - 3) + (x^4 + 3x^3 - 8)$

i $3(7 + 4x - x^2 - 2x^3) + 5(-2 - 3x + x^3)$

j $6(x^3 + 5x^2 - 2) - 3(2x^3 - x^2 - x)$

k $8(x^4 + 2x^2 - 4x - 1) - 2(5 - 3x + x^3)$

l $7(x^6 + 3x^3 + x^2 - 4) - 4(2x^6 + x^5 - 3x - 7)$

2 Simplify

a $(3y^2 + 2y + 1) + (y^3 - 4y^2 + 7y) + (2y^3 - y^2 - 8y + 5)$

b $3(t^4 - t^3 + 4t) + (6 - t - 3t^3) + 2(t^4 - 2t^2 + 4)$

c $(x^3 - 6x^2 + 8) + (5x^2 - x + 1) - (2x^3 + 3x^2 + x - 7)$

d $2(3 + m + 7m^2 - 3m^5) + 6(1 - m^2 + 2m^4) - 5(m^5 + 3m^3 - m^2 + 2)$

e $\frac{1}{3}(1 - 2u + \frac{3}{5}u^2 + 3u^4) - \frac{1}{2}(2 - u + \frac{2}{3}u^2 - \frac{1}{2}u^3)$

3 Giving your answers in ascending powers of x , simplify

a $x(2 - 3x + x^2) + 4(1 + 2x^2 - x^3)$

b $x(x^4 + 7x^2 - 5x + 9) - 2(x^4 - 4x^3 - 3)$

c $2x(-5 + 4x - x^3) + 7(2 - 3x^2 + x^4)$

d $x^2(8 + 2x + x^2) - 3(5 + 4x^2 + x^3)$

e $3x^2(x + 3) - x(x^3 + 4x^2) + 5(x^3 - 2x)$

f $x^2(6 - x + 5x^2) + 7x(2 - x^3) + 4(1 - 3x - x^2)$

4 Show that

a $(3x + 1)(x^2 - 2x + 4) \equiv 3x^3 - 5x^2 + 10x + 4$

b $(1 + 2x - x^2)(1 - 2x + x^2) \equiv 1 - 4x^2 + 4x^3 - x^4$

c $(3 - x)^3 \equiv 27 - 27x + 9x^2 - x^3$

5 Giving your answers in descending powers of x , expand and simplify

a $(x + 1)(x^2 + 5x - 6)$

b $(2x - 5)(x^2 - 3x + 7)$

c $(4 - 7x)(2 + 5x - x^2)$

d $(3x - 2)^3$

e $(x^2 + 3)(2x^2 - x + 9)$

f $(4x - 1)(x^4 - 3x^2 + 5x + 2)$

g $(x^2 + 2x + 5)(x^2 + 3x + 1)$

h $(x^2 + x - 3)(2x^2 - x + 4)$

i $(3x^2 - 5x + 2)(2x^2 - 4x - 8)$

j $(x^2 + 2x - 6)^2$

k $(x^3 + 4x^2 + 1)(2x^4 + x^2 + 3)$

l $(6 - 2x + x^3)(3 + x^2 - x^3 + 2x^4)$

6 Simplify

a $(p^2 - 1)(p + 4)(2p + 3)$

b $(t + 2)(t^2 + 3t + 5) + (t + 4)(t^2 + t + 7)$

c $2(x^2 - 3)(x^2 + x - 4) + (3x - 1)(4x^3 + 2x^2 - x + 6)$

d $(u^3 - 4u^2 - 3)(u + 2) - (2u^3 + u - 1)(u^2 + 5u - 3)$

1 Factorise

a $x^2 + 4x + 3$

b $x^2 + 7x + 10$

c $y^2 - 3y + 2$

d $x^2 - 6x + 9$

e $y^2 - y - 2$

f $a^2 + 2a - 8$

g $x^2 - 1$

h $p^2 + 9p + 14$

i $x^2 - 2x - 15$

j $16 - 10m + m^2$

k $t^2 + 3t - 18$

l $y^2 - 13y + 40$

m $r^2 - 16$

n $y^2 - 2y - 63$

o $121 + 22a + a^2$

p $x^2 + 6x - 72$

q $26 - 15x + x^2$

r $s^2 + 23s + 120$

s $p^2 + 14p - 51$

t $m^2 - m - 90$

2 Factorise

a $2x^2 + 3x + 1$

b $2 + 7p + 3p^2$

c $2y^2 - 5y + 3$

d $2 - m - m^2$

e $3r^2 - 2r - 1$

f $5 - 19y - 4y^2$

g $4 - 13a + 3a^2$

h $5x^2 - 8x - 4$

i $4x^2 + 8x + 3$

j $9s^2 - 6s + 1$

k $4m^2 - 25$

l $2 - y - 6y^2$

m $4u^2 + 17u + 4$

n $6p^2 + 5p - 4$

o $8x^2 + 19x + 6$

p $12r^2 + 8r - 15$

3 Using factorisation, solve each equation.

a $x^2 - 4x + 3 = 0$

b $x^2 + 6x + 8 = 0$

c $x^2 + 4x - 5 = 0$

d $x^2 - 7x = 8$

e $x^2 - 25 = 0$

f $x(x - 1) = 42$

g $x^2 = 3x$

h $27 + 12x + x^2 = 0$

i $60 - 4x - x^2 = 0$

j $5x + 14 = x^2$

k $2x^2 - 3x + 1 = 0$

l $x(x - 1) = 6(x - 2)$

m $3x^2 + 11x = 4$

n $x(2x - 3) = 5$

o $6 + 23x - 4x^2 = 0$

p $6x^2 + 10 = 19x$

q $4x^2 + 4x + 1 = 0$

r $3(x^2 + 4) = 13x$

s $(2x + 5)^2 = 5 - x$

t $3x(2x - 7) = 2(7x + 3)$

4 Factorise fully

a $2y^2 - 10y + 12$

b $x^3 + x^2 - 2x$

c $p^3 - 4p$

d $3m^3 + 21m^2 + 18m$

e $a^4 + 4a^2 + 3$

f $t^4 + 3t^2 - 10$

g $12 + 20x - 8x^2$

h $6r^2 - 9r - 42$

i $6x^3 - 26x^2 + 8x$

j $y^4 + 3y^3 - 18y^2$

k $m^4 - 1$

l $p^5 - 4p^3 + 4p$

5 Sketch each curve showing the coordinates of any points of intersection with the coordinate axes.

a $y = x^2 - 3x + 2$

b $y = x^2 + 5x + 6$

c $y = x^2 - 9$

d $y = x^2 - 2x$

e $y = x^2 - 10x + 25$

f $y = 2x^2 - 14x + 20$

g $y = -x^2 + 5x - 4$

h $y = 2 + x - x^2$

i $y = 2x^2 - 3x + 1$

j $y = 2x^2 + 13x + 6$

k $y = 3 - 8x + 4x^2$

l $y = 2 + 7x - 4x^2$

m $y = 5x^2 - 17x + 6$

n $y = -6x^2 + 7x - 2$

o $y = 6x^2 + x - 5$

6 Solve each of the following equations.

a $x - 5 + \frac{4}{x} = 0$

b $x - \frac{10}{x} = 3$

c $2x^3 - x^2 - 3x = 0$

d $x^2(10 - x^2) = 9$

e $\frac{5}{x^2} + \frac{4}{x} - 1 = 0$

f $\frac{x-6}{x-4} = x$

g $x + 5 = \frac{3}{x+3}$

h $x^2 - \frac{4}{x^2} = 3$

i $4x^4 + 7x^2 = 2$

j $\frac{2x}{3-x} = \frac{1}{x+2}$

k $\frac{2x+1}{x+3} = \frac{2}{x}$

l $\frac{7}{x+2} - 3x = 2$

- 1 Express in the form $(x + a)^2 + b$
- | | | | |
|----------------------------|----------------------------------|-----------------------------------|---|
| a $x^2 + 2x + 4$ | b $x^2 - 2x + 4$ | c $x^2 - 4x + 1$ | d $x^2 + 6x$ |
| e $x^2 + 4x + 8$ | f $x^2 - 8x - 5$ | g $x^2 + 12x + 30$ | h $x^2 - 10x + 25$ |
| i $x^2 + 6x - 9$ | j $18 - 4x + x^2$ | k $x^2 + 3x + 3$ | l $x^2 + x - 1$ |
| m $x^2 - 18x + 100$ | n $x^2 - x - \frac{1}{2}$ | o $20 + 9x + x^2$ | p $x^2 - 7x - 2$ |
| q $5 - 3x + x^2$ | r $x^2 - 11x + 37$ | s $x^2 + \frac{2}{3}x + 1$ | t $x^2 - \frac{1}{2}x - \frac{1}{4}$ |
- 2 Express in the form $a(x + b)^2 + c$
- | | | | |
|----------------------------|---------------------------|----------------------------|--|
| a $2x^2 + 4x + 3$ | b $2x^2 - 8x - 7$ | c $3 - 6x + 3x^2$ | d $4x^2 + 24x + 11$ |
| e $-x^2 - 2x - 5$ | f $1 + 10x - x^2$ | g $2x^2 + 2x - 1$ | h $3x^2 - 9x + 5$ |
| i $3x^2 - 24x + 48$ | j $3x^2 - 15x$ | k $70 + 40x + 5x^2$ | l $2x^2 + 5x + 2$ |
| m $4x^2 + 6x - 7$ | n $-2x^2 + 4x - 1$ | o $4 - 2x - 3x^2$ | p $\frac{1}{3}x^2 + \frac{1}{2}x - \frac{1}{4}$ |
- 3 Solve each equation by completing the square, giving your answers as simply as possible in terms of surds where appropriate.
- | | | | |
|-----------------------------|-------------------------------|-----------------------------|------------------------------|
| a $y^2 - 4y + 2 = 0$ | b $p^2 + 2p - 2 = 0$ | c $x^2 - 6x + 4 = 0$ | d $7 + 10r + r^2 = 0$ |
| e $x^2 - 2x = 11$ | f $a^2 - 12a - 18 = 0$ | g $m^2 - 3m + 1 = 0$ | h $9 - 7t + t^2 = 0$ |
| i $u^2 + 7u = 44$ | j $2y^2 - 4y + 1 = 0$ | k $3p^2 + 18p = -23$ | l $2x^2 + 12x = 9$ |
| m $-m^2 + m + 1 = 0$ | n $4x^2 + 49 = 28x$ | o $1 - t - 3t^2 = 0$ | p $2a^2 - 7a + 4 = 0$ |
- 4 By completing the square, find the maximum or minimum value of y and the value of x for which this occurs. State whether your value of y is a maximum or a minimum in each case.
- | | | |
|-------------------------------|-------------------------------|-------------------------------|
| a $y = x^2 - 2x + 7$ | b $y = x^2 + 2x - 3$ | c $y = 1 - 6x + x^2$ |
| d $y = x^2 + 10x + 35$ | e $y = -x^2 + 4x + 4$ | f $y = x^2 + 3x - 2$ |
| g $y = 2x^2 + 8x + 5$ | h $y = -3x^2 + 6x$ | i $y = 7 - 5x - x^2$ |
| j $y = 4x^2 - 12x + 9$ | k $y = 4x^2 + 20x - 8$ | l $y = 17 - 2x - 2x^2$ |
- 5 Sketch each curve showing the exact coordinates of its turning point and the point where it crosses the y -axis.
- | | | |
|------------------------------|--------------------------------|--------------------------------|
| a $y = x^2 - 4x + 3$ | b $y = x^2 + 2x - 24$ | c $y = x^2 - 2x + 5$ |
| d $y = 30 + 8x + x^2$ | e $y = x^2 + 2x + 1$ | f $y = 8 + 2x - x^2$ |
| g $y = -x^2 + 8x - 7$ | h $y = -x^2 - 4x - 7$ | i $y = x^2 - 5x + 4$ |
| j $y = x^2 + 3x + 3$ | k $y = 3 + 8x + 4x^2$ | l $y = -2x^2 + 8x - 15$ |
| m $y = 1 - x - 2x^2$ | n $y = 25 - 20x + 4x^2$ | o $y = 3x^2 - 4x + 2$ |
- 6 **a** Express $x^2 - 4\sqrt{2}x + 5$ in the form $a(x + b)^2 + c$.
b Write down an equation of the line of symmetry of the curve $y = x^2 + 4\sqrt{2}x + 5$.
- 7 $f(x) \equiv x^2 + 2kx - 3$.
- By completing the square, find the roots of the equation $f(x) = 0$ in terms of the constant k .


- 1 By completing the square, show that the roots of the equation $ax^2 + bx + c = 0$ are given by

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

- 2 Use the quadratic formula to solve each equation, giving your answers as simply as possible in terms of surds where appropriate.

a $x^2 + 4x + 1 = 0$	b $4 + 8t - t^2 = 0$	c $y^2 - 20y + 91 = 0$	d $r^2 + 2r - 7 = 0$
e $6 + 18a + a^2 = 0$	f $m(m - 5) = 5$	g $x^2 + 11x + 27 = 0$	h $2u^2 + 6u + 3 = 0$
i $5 - y - y^2 = 0$	j $2x^2 - 3x = 2$	k $3p^2 + 7p + 1 = 0$	l $t^2 - 14t = 14$
m $0.1r^2 + 1.4r = 0.9$	n $6u^2 + 4u = 1$	o $\frac{1}{2}y^2 - 3y = \frac{2}{3}$	p $4x(x - 3) = 11 - 4x$

- 3



The diagram shows the curve with equation $y = 2x^2 - 8x + 3$.

Find and simplify the exact coordinates of the points where the curve crosses the x -axis.

- 4 State the condition for which the roots of the equation $ax^2 + bx + c = 0$ are

a real and distinct **b** real and equal **c** not real

- 5 Sketch the curve $y = ax^2 + bx + c$ and the x -axis in the cases where

a $a > 0$ and $b^2 - 4ac > 0$	b $a < 0$ and $b^2 - 4ac < 0$
c $a > 0$ and $b^2 - 4ac = 0$	d $a < 0$ and $b^2 - 4ac > 0$

- 6 By evaluating the discriminant, determine whether the roots of each equation are real and distinct, real and equal or not real.

a $x^2 + 2x - 7 = 0$	b $x^2 + x + 3 = 0$	c $x^2 - 4x + 5 = 0$	d $x^2 - 6x + 3 = 0$
e $x^2 + 14x + 49 = 0$	f $x^2 - 9x + 17 = 0$	g $x^2 + 3x = 11$	h $2 + 3x + 2x^2 = 0$
i $5x^2 + 8x + 3 = 0$	j $3x^2 - 7x + 5 = 0$	k $9x^2 - 12x + 4 = 0$	l $13x^2 + 19x + 7 = 0$
m $4 - 11x + 8x^2 = 0$	n $x^2 + \frac{2}{3}x = \frac{1}{4}$	o $x^2 - \frac{3}{4}x + \frac{1}{8} = 0$	p $\frac{2}{5}x^2 + \frac{3}{5}x + \frac{1}{3} = 0$

- 7 Find the value of the constant p such that the equation $x^2 + x + p = 0$ has equal roots.

- 8 Given that $q \neq 0$, find the value of the constant q such that the equation $x^2 + 2qx - q = 0$ has a repeated root.

- 9 Given that the x -axis is a tangent to the curve with the equation

$$y = x^2 + rx - 2x + 4,$$

find the two possible values of the constant r .

- 1 a Factorise fully the expression

$$20x - 2x^2 - 6x^3.$$

- b Hence, find all solutions to the equation

$$20x - 2x^2 - 6x^3 = 0.$$

- 2 A is the point $(-2, 1)$ and B is the point $(6, k)$.

- a Show that $AB^2 = k^2 - 2k + 65$.

Given also that $AB = 10$,

- b find the possible values of k .

- 3 Solve the equations

a $x - \frac{5}{x} = 4$

b $\frac{9}{5-x} - 1 = 2x$


- 4 a Find the coordinates of the turning point of the curve with equation $y = 3 - 5x - 2x^2$.

- b Sketch the curve $y = 3 - 5x - 2x^2$, showing the coordinates of any points of intersection with the coordinate axes.

- 5 Find in the form $k\sqrt{2}$ the solutions of the equation

$$2x^2 + 5\sqrt{2}x - 6 = 0.$$

- 6



The diagram shows the curve with equation $y = 3x^2 - 9x + k$ where k is a constant.

- a Find the x -coordinate of the turning point of the curve, P .

Given that the y -coordinate of P is $\frac{17}{4}$,

- b find the coordinates of the point Q where the curve crosses the y -axis.

- 7 By letting $y = 2^x$, or otherwise, solve the equation

$$2^{2x} - 10(2^x) + 16 = 0.$$

- 8 Given that the equation

$$kx^2 - 2x + 3 - 2k = 0$$

has equal roots, find the possible values of the constant k .

- 9 $f(x) \equiv 3 + 4x - x^2$.
- a Express $f(x)$ in the form $a(x + b)^2 + c$.
- b State the coordinates of the turning point of the curve $y = f(x)$.
- c Solve the equation $f(x) = 2$, giving your answers in the form $d + e\sqrt{5}$.
- 10 Giving your answers in terms of surds, solve the equations
- a $3x^2 - 5x + 1 = 0$
- b $\frac{x}{x+2} = \frac{3}{x-1}$
- 11 a By completing the square, find, in terms of k , the solutions of the equation
 $x^2 - 4kx + 6 = 0$.
- b Using your answers to part a, solve the equation
 $x^2 - 12x + 6 = 0$.
- 12 a Find in the form $a + b\sqrt{3}$, where a and b are integers, the values of x such that
 $2x^2 - 12x = 6$.
- b Solve the equation
 $2y^3 + y^2 - 15y = 0$.
- 13 Labelling the coordinates of any points of intersection with the coordinate axes, sketch the curves
- a $y = (x + 1)(x - p)$ where $p > 0$,
- b $y = (x + q)^2$ where $q < 0$.
- 14 $f(x) \equiv 2x^2 - 6x + 5$.
- a Find the values of A , B and C such that
 $f(x) \equiv A(x + B)^2 + C$.
- b Hence deduce the minimum value of $f(x)$.
- 15 a Given that $t = x^{\frac{1}{3}}$ express $x^{\frac{2}{3}}$ in terms of t .
- b Hence, or otherwise, solve the equation
 $2x^{\frac{2}{3}} + x^{\frac{1}{3}} - 6 = 0$.
- 16 a Express $k^2 - 8k + 20$ in the form $a(k + b)^2 + c$, where a , b and c are constants.
- b Hence prove that the equation
 $x^2 - kx + 2k = 5$
has real and distinct roots for all real values of k .
- 17 a Show that
 $(x^2 + 2x - 3)(x^2 - 3x - 4) \equiv x^4 - x^3 - 13x^2 + x + 12$.
- b Hence solve the equation
 $x^4 - x^3 - 13x^2 + x + 12 = 0$.

1 Solve each pair of simultaneous equations.

a $y = 3x$

$y = 2x + 1$

b $y = x - 6$

$y = \frac{1}{2}x - 4$

c $y = 2x + 6$

$y = 3 - 4x$

d $x + y - 3 = 0$

$x + 2y + 1 = 0$

e $x + 2y + 11 = 0$

$2x - 3y + 1 = 0$

f $3x + 3y + 4 = 0$

$5x - 2y - 5 = 0$

2 Find the coordinates of the points of intersection of the given straight line and curve in each case.

a $y = x + 2$

$y = x^2 - 4$

b $y = 4x + 11$

$y = x^2 + 3x - 1$

c $y = 2x - 1$

$y = 2x^2 + 3x - 7$

3 Solve each pair of simultaneous equations.

a $x^2 - y + 3 = 0$

$x - y + 5 = 0$

b $2x^2 - y - 8x = 0$

$x + y + 3 = 0$

c $x^2 + y^2 = 25$

$2x - y = 5$

d $x^2 + 2xy + 15 = 0$

$2x - y + 10 = 0$

e $x^2 - 2xy - y^2 = 7$

$x + y = 1$

f $3x^2 - x - y^2 = 0$

$x + y - 1 = 0$

g $2x^2 + xy + y^2 = 22$

$x + y = 4$

h $x^2 - 4y - y^2 = 0$

$x - 2y = 0$

i $x^2 + xy = 4$

$3x + 2y = 6$

j $2x^2 + y - y^2 = 8$

$2x - y = 3$

k $x^2 - xy + y^2 = 13$

$2x - y = 7$

l $x^2 - 5x + y^2 = 0$

$3x + y = 5$

m $3x^2 - xy + y^2 = 36$

$x - 2y = 10$

n $2x^2 + x - 4y = 6$

$3x - 2y = 4$

o $x^2 + x + 2y^2 - 52 = 0$

$x - 3y + 17 = 0$

4 Solve each pair of simultaneous equations.

a $x - \frac{1}{y} - 4y = 0$

$x - 6y - 1 = 0$

b $xy = 6$

$x - y = 5$

c $\frac{3}{x} - 2y + 4 = 0$

$4x + y - 7 = 0$

5 The line $y = 5 - x$ intersects the curve $y = x^2 - 3x + 2$ at the points P and Q .

Find the length PQ in the form $k\sqrt{2}$.

6 Solve the simultaneous equations

$$3^{x-1} = 9^{2y}$$

$$8^{x-2} = 4^{1+y}$$

7 Given that

$$(A + 2\sqrt{3})(B - \sqrt{3}) \equiv 9\sqrt{3} - 1,$$

find the values of the integers A and B .

1 Find the set of values of x for which

- a** $2x + 1 < 7$ **b** $3x - 1 \geq 20$ **c** $2x - 5 > 3$ **d** $6 + 3x \leq 42$
e $5x + 17 \geq 2$ **f** $\frac{1}{3}x + 7 < 8$ **g** $9x - 4 \geq 50$ **h** $3x + 11 < 7$
i $18 - x > 4$ **j** $10 + 4x \leq 0$ **k** $12 - 3x < 10$ **l** $9 - \frac{1}{2}x \geq 4$

2 Solve each inequality.

- a** $2y - 3 > y + 4$ **b** $5p + 1 \leq p + 3$ **c** $x - 2 < 3x - 8$
d $a + 11 \geq 15 - a$ **e** $17 - 2u < 2 + u$ **f** $5 - b \geq 14 - 3b$
g $4x + 23 < x + 5$ **h** $12 + 3y \geq 2y - 1$ **i** $16 - 3p \leq 36 + p$
j $5(r - 2) > 30$ **k** $3(1 - 2t) \leq t - 4$ **l** $2(3 + x) \geq 4(6 - x)$
m $7(y + 3) - 2(3y - 1) < 0$ **n** $4(5 - 2x) > 3(7 - 2x)$ **o** $3(4u - 1) - 5(u - 3) < 9$

3 Find the set of values of x for which

- a** $x^2 - 4x + 3 < 0$ **b** $x^2 - 4 \leq 0$ **c** $15 + 8x + x^2 < 0$ **d** $x^2 + 2x \leq 8$
e $x^2 - 6x + 5 > 0$ **f** $x^2 + 4x > 12$ **g** $x^2 + 10x + 21 \geq 0$ **h** $22 + 9x - x^2 > 0$
i $63 - 2x - x^2 \leq 0$ **j** $x^2 + 11x + 30 > 0$ **k** $30 + 7x - x^2 > 0$ **l** $x^2 + 91 \geq 20x$

4 Solve each inequality.

- a** $2x^2 - 9x + 4 \leq 0$ **b** $2r^2 - 5r - 3 < 0$ **c** $2 - p - 3p^2 \geq 0$
d $2y^2 + 9y - 5 > 0$ **e** $4m^2 + 13m + 3 < 0$ **f** $9x - 2x^2 \leq 10$
g $a^2 + 6 < 8a - 9$ **h** $x(x + 4) \leq 7 - 2x$ **i** $y(y + 9) > 2(y - 5)$
j $x(2x + 1) > x^2 + 6$ **k** $u(5 - 6u) < 3 - 4u$ **l** $2t + 3 \geq 3t(t - 2)$
m $(y - 2)^2 \leq 2y - 1$ **n** $(p + 2)(p + 3) \geq 20$ **o** $2(13 + 2x) < (6 + x)(1 - x)$

5 Giving your answers in terms of surds, find the set of values of x for which

- a** $x^2 + 2x - 1 < 0$ **b** $x^2 - 6x + 4 > 0$ **c** $11 - 6x - x^2 > 0$ **d** $x^2 + 4x + 1 \geq 0$

6 Find the value or set of values of k such that

- a** the equation $x^2 - 6x + k = 0$ has equal roots,
b the equation $x^2 + 2x + k = 0$ has real and distinct roots,
c the equation $x^2 - 3x + k = 0$ has no real roots,
d the equation $x^2 + kx + 4 = 0$ has real roots,
e the equation $kx^2 + x - 1 = 0$ has equal roots,
f the equation $x^2 + kx - 3k = 0$ has no real roots,
g the equation $x^2 + 2x + k - 2 = 0$ has real and distinct roots,
h the equation $2x^2 - kx + k = 0$ has equal roots,
i the equation $x^2 + kx + 2k - 3 = 0$ has no real roots,
j the equation $3x^2 + kx - x + 3 = 0$ has real roots.

1 Solve each of the following inequalities.


a $\frac{1}{2}y + 3 > 2y - 1$

b $x^2 - 8x + 12 \geq 0$

2 Find the set of integers, n , for which

$$2n^2 - 5n < 12.$$

3



The diagram shows a rectangular birthday card which is x cm wide and $(x + 8)$ cm tall.

Given that the height of the card is to be at least 50% more than its width,

a show that $x \leq 16$.

Given also that the area of the front of the card is to be at least 180 cm^2 ,

b find the set of possible values of x .

4 Find the set of values of x for which

$$(3x - 1)^2 < 5x - 1.$$

5 Given that $x - y = 8$,

and that $xy \leq 240$,

find the maximum value of $(x + y)$.

6 Solve the inequality


$$(3t + 1)(t - 4) \geq 2t(t - 7).$$

7 Given that the equation $2x(x + 1) = kx - 8$ has real and distinct roots,

a show that $k^2 - 4k - 60 > 0$,

b find the set of possible values of k .

8



A party hat is designed in the shape of a right circular cone of base radius r cm and slant height $(3r - 4)$ cm.

Given that the height of the cone must not be more than 24 cm, find the maximum value of r .

- 1 a Find the value of x such that
$$2^{x-1} = 16. \quad (3)$$
- b Find the value of y such that
$$2(3^y - 10) = 34. \quad (2)$$
- 2 a Express $x^2 - 6x + 11$ in the form $(x + a)^2 + b$. (2)
- b Sketch the curve $y = x^2 - 6x + 11$, and show the coordinates of the turning point of the curve. (3)
- 3 a Express $(12\frac{1}{4})^{-\frac{1}{2}}$ as an exact fraction in its simplest form. (2)
- b Solve the equation
$$3x^{-3} = 7\frac{1}{9}. \quad (3)$$
- 4 Solve the equation
$$x\sqrt{12} + 9 = x\sqrt{3},$$

giving your answer in the form $k\sqrt{3}$, where k is an integer. (4)
- 5 a Solve the equation
$$x^2 + 10x + 13 = 0,$$

giving your answers in the form $a + b\sqrt{3}$, where a and b are integers. (4)
- b Hence find the set of values of x for which
$$x^2 + 10x + 13 > 0. \quad (2)$$
- 6 Solve the equations
- a $7(6x - 7) = 9x^2$ (3)
- b $\frac{2}{y+1} + 1 = 2y$ (4)
- 7 Solve the simultaneous equations
$$x - y + 3 = 0$$

$$3x^2 - 2xy + y^2 - 17 = 0 \quad (6)$$
- 8 a Find the value of x such that
$$x^{\frac{3}{2}} = 64. \quad (2)$$
- b Given that
$$\frac{\sqrt{3}+1}{2\sqrt{3}-3} \equiv a + b\sqrt{3},$$

find the values of the rational constants a and b . (4)
- 9 The point $P(2k, k)$ lies within a circle of radius 3, centre $(2, 4)$.
- a Show that $5k^2 - 16k + 11 < 0$. (4)
- b Hence find the set of possible values of k . (3)

10 Solve each of the following inequalities.

a $4x - 1 \leq 2x + 6$ (2)

b $x(2x + 1) < 1$ (4)

11 $f(x) = 2x^2 - 8x + 5$.

a Express $f(x)$ in the form $a(x + b)^2 + c$, where a , b and c are integers. (3)

b Write down the coordinates of the turning point of the curve $y = f(x)$. (1)

c Solve the equation $f(x) = 0$, giving your answers in the form $p + q\sqrt{6}$, where p and q are rational. (3)

12 Simplify

a $\sqrt{12} - \frac{5}{\sqrt{3}}$ (3)


b $\frac{(4\sqrt{x})^3}{16x}$ (2)

13 Given that the equation

$$x^2 - 2kx + k + 6 = 0$$

has no real roots, find the set of possible values of the constant k . (6)

14



The diagram shows triangle ABC in which $AB = BC = 4 + \sqrt{3}$ and $AC = 4 + 4\sqrt{3}$.

Given that M is the mid-point of AC ,

a find the exact length BM , (4)

b show that the area of triangle ABC is $6 + 2\sqrt{3}$. (2)

15 Solve the equation

$$4^{2y+7} = 8^{y+3}. \quad (4)$$

16 Show that

$$(x^2 - x + 3)(2x^2 - 3x - 9) \equiv Ax^4 + Bx^3 + C,$$

where A , B and C are constants to be found. (4)

17 $f(x) = x^2 + 4x + k$.

a By completing the square, find in terms of the constant k the roots of the equation $f(x) = 0$. (4)

b State the set of values of k for which the equation $f(x) = 0$ has real roots. (1)

c Use your answers to part a to solve the equation

$$x^2 + 4x - 4 = 0,$$

giving your answers in the form $a + b\sqrt{2}$, where a and b are integers. (2)

- 1 Solve the inequality

$$(x + 1)(x + 2) \leq 12. \quad (5)$$


- 2 a Express $2^{\frac{7}{2}} - 2^{\frac{3}{2}}$ in the form $k\sqrt{2}$. (2)

- b Show that

$$(\sqrt{x} + 6)^2 + (2\sqrt{x} - 3)^2$$

can be written in the form $ax + b$ where a and b are integers to be found. (3)

- 3



The diagram shows the curve with equation $y = 2x^2 + px + q$, where p and q are constants, which crosses the x -axis at the points with coordinates $(-2, 0)$ and $(3, 0)$.

- a Show that $p = -2$ and find the value of q . (4)

- b Find the coordinates of the turning point of the curve. (3)

- 4 Solve the equation

$$2(x - \sqrt{32}) = \sqrt{98} - x,$$

giving your answer in the form $k\sqrt{2}$. (4)

- 5 Given that the equation

$$kx^2 - 4kx + 3 = 0,$$

where k is a constant, has real and distinct roots,

- a show that $k(4k - 3) > 0$, (3)

- b find the set of possible values of k . (2)

- 6 Solve the simultaneous equations

$$4^{2x} = 2^{y-1}$$

$$9^{4x} = 3^{y+1} \quad (7)$$

- 7 a Find the values of the constants a and b such that

$$x^2 - 7x + 9 \equiv (x + a)^2 + b. \quad (3)$$

- b Hence, write down an equation of the line of symmetry of the curve $y = x^2 - 7x + 9$. (1)

- 8 a Solve the inequality

$$y^2 - 2y - 15 < 0. \quad (3)$$

- b Find the exact values of x for which

$$\frac{x}{x-3} = \frac{4}{2-x}. \quad (5)$$

- 9 Solve the equation

$$2^{x^2+2} = 8^x. \quad (5)$$

- 10 Giving your answers in terms of surds, solve the equations

a $t(1 - 2t) = 3(t - 5)$ (4)

b $x^4 - x^2 - 6 = 0$ (4)

- 11 Find the set of values of x for which

$$21 - 4x - x^2 \leq 0. \quad (4)$$

- 12 a Given that $y = 3^x$ express 3^{2x+2} in terms of y . (2)

- b Hence, or otherwise, solve the equation

$$3^{2x+2} - 10(3^x) + 1 = 0. \quad (4)$$

- 13 a Express $5\sqrt{3}$ in the form \sqrt{k} . (2)

- b Hence find the integer n such that

$$n < 5\sqrt{3} < n + 1. \quad (3)$$

- 14 Solve the simultaneous equations

$$\begin{aligned} 2x^2 - y^2 - 7 &= 0 \\ 2x - 3y + 7 &= 0 \end{aligned} \quad (8)$$

- 15 Express each of the following in the form $a + b\sqrt{2}$, where a and b are integers.

a $\frac{\sqrt{48} - \sqrt{600}}{\sqrt{12}}$ (3)

b $\frac{\sqrt{2}}{4 + 3\sqrt{2}}$ (4)

- 16 Given that $5^{x+1} = 25^{y-3}$,

- a find an expression for y in terms of x . (4)

Given also that $16^{x-1} = 4^z$,

- b find an expression for z in terms of y . (4)

- 17 a By completing the square, find in terms of the constant k the roots of the equation

$$x^2 - 2kx - k = 0. \quad (4)$$

- b Hence, find the set of values of k for which the equation has real roots. (3)

- 18 a Given that $y = x^{\frac{1}{5}}$, show that the equation

$$x^{-\frac{1}{5}} - x^{\frac{1}{5}} = \frac{3}{2}$$

can be written as

$$2y^2 + 3y - 2 = 0. \quad (3)$$

- b Hence find the values of x for which

$$x^{-\frac{1}{5}} - x^{\frac{1}{5}} = \frac{3}{2}. \quad (4)$$

1 a Express $(\frac{2}{3})^{-2}$ as an exact fraction in its simplest form. (2)

b Solve the equation

$$x^{\frac{3}{2}} - 27 = 0. \quad (3)$$

2 Solve the simultaneous equations

$$x + 3y = 16$$

$$x^2 - xy + 2y^2 = 46 \quad (7)$$

3 Simplify

a $\sqrt{192} - 2\sqrt{12} + \sqrt{75}$ (4)


b $(2 + \sqrt{3})(5 - 2\sqrt{3})$ (3)

4 $f(x) \equiv x^2 - 4\sqrt{2}x + 11.$

a Express $f(x)$ in the form $a(x + b)^2 + c$ stating the exact values of the constants a , b and c . (4)

b Sketch the curve $y = f(x)$, showing the coordinates of the turning point of the curve and of any points of intersection of the curve with the coordinate axes. (3)

5



A sealed metal container for food is a cylinder of height 12 cm and base radius r cm.

Given that the surface area of the container must be at most $128\pi \text{ cm}^2$,

a show that $r^2 + 12r - 64 \leq 0$. (3)

b Hence find the maximum value of r . (4)

6 Find the non-zero value of x for which

$$(2\sqrt{x})^3 = 4x. \quad (4)$$

7 a Write down the value of x such that $2^x = 32$. (1)

b Solve the equation

$$32^{y+1} = 4^y. \quad (3)$$

8 a Given that $t = \sqrt{x}$, express $x - 5\sqrt{x}$ in terms of t . (1)

b Hence, or otherwise, solve the equation

$$x - 5\sqrt{x} + 6 = 0. \quad (4)$$

9 Prove, by completing the square, that there is no real value of the constant k for which the equation $x^2 + kx + 3 + k^2 = 0$ has real roots. (6)

- 10 a Find the value of x such that

$$8^{2x-1} = 32. \quad (3)$$
- b Find the value of y such that

$$\left(\frac{1}{3}\right)^{y-2} = 81. \quad (3)$$
- 11 Solve the inequality

$$x(2x - 7) < (x - 2)^2. \quad (5)$$
- 12 Express

$$\frac{2}{3\sqrt{2}-4} - \frac{3-\sqrt{2}}{\sqrt{2}+1}$$
in the form $a + b\sqrt{2}$, where a and b are integers. (6)
- 13 a Solve the equation

$$6y^2 + 25y - 9 = 0. \quad (3)$$
- b Find the values of the constant k such that the equation

$$x^2 + kx + 16 = 0$$
has equal roots. (3)
- 14 a Given that $y = 2^x$,
i show that $4^x = y^2$,
ii express 2^{x-1} in terms of y . (4)
- b By using your answers to part a, or otherwise, find the values of x for which

$$4^x - 9(2^{x-1}) + 2 = 0. \quad (4)$$
- 15 Find the pairs of values (x, y) which satisfy the simultaneous equations

$$x^2 + 2xy + y^2 = 9$$

$$x - 3y = 1 \quad (7)$$
- 16 a Prove, by completing the square, that the roots of the equation $x^2 + ax + b = 0$ are given by

$$x = \frac{-a \pm \sqrt{a^2 - 4b}}{2}. \quad (6)$$
- b Hence, find an expression for b in terms of a such that the equation $x^2 + ax + b = 0$ has a repeated root. (2)
- 17
$$f(x) \equiv 2x^2 - 12x + 19.$$
- a Prove that $f(x) \geq 1$ for all real values of x . (5)
- b Find the set of values of x for which $f(x) < 9$. (4)
- 18 a Express $(1 - \sqrt{5})^2$ in the form $a + b\sqrt{5}$. (2)
- b Hence, or otherwise, solve the equation

$$y^2 = 3 - \sqrt{5},$$
giving your answers in the form $c\sqrt{2} + d\sqrt{10}$, where c and d are exact fractions. (6)