


2



Review Exercise

- 1 a** On the same set of axes sketch the graphs of $y = |2x - 1|$ and $y = |x - k|$, $k > 1$.
- b** Find, in terms of k , the values of x for which $|2x - 1| = |x - k|$.

- 2 a** Sketch the graph of $y = |3x + 2| - 4$, showing the coordinates of the points of intersection of the graph with the axes.
- b** Find the values of x for which $|3x + 2| = 4 + x$.




The figure shows the graph of $y = f(x)$, $-5 \leq x \leq 5$.

The point $M(2, 4)$ is the maximum turning point of the graph.

Sketch, on separate diagrams, the graphs of

- a** $y = f(x) + 3$ **b** $y = |f(x)|$
c $y = f(|x|)$.

Show on each graph the coordinates of any maximum turning points. **E**



The diagram shows a sketch of the graph of the increasing function f .

The curve crosses the x -axis at the point $A(1, 0)$ and the y -axis at the point $B(0, -2)$.

On separate diagrams, sketch the graph of:

- a** $y = f^{-1}(x)$ **b** $y = f(|x|)$
c $y = f(2x) + 1$ **d** $y = 3f(x - 1)$.

In each case, show the images of the points A and B .

- 5** For the positive constant k , where $k > 1$, the functions f and g are defined by
- $f: x \rightarrow \ln(x + k), x > -k,$
 $g: x \rightarrow |2x - k|, x \in \mathbb{R}$

a Sketch, on the same set of axes, the graphs of f and g . Give the coordinates of points where the graphs meet the axes.

b Write down the range of f .


c Find, in terms of k , $fg\left(\frac{k}{4}\right)$.

The curve C has equation $y = f(x)$. The tangent to C at the point with x -coordinate 3 is parallel to the line with equation $9y = 2x + 1$.

d Find the value of k .

E

6



The diagram shows a sketch of the graph of $y = f(x)$.

The curve has a minimum at the point $A(1, -1)$, passes through x -axis at the origin, and the points $B(2, 0)$ and $C(5, 0)$; the asymptotes have equations $x = 3$ and $y = 2$.

a Sketch, on separate axes, the graph of

i $y = |f(x)|$

ii $y = -f(x + 1)$


iii $y = f(-2x)$

in each case, showing the images of the points A , B and C .

b State the number of solutions to the equation

i $3|f(x)| = 2$ **ii** $2|f(x)| = 3$.

7



The diagram shows part of the curve C with equation $y = f(x)$ where

$$f(x) = \frac{(x - 1)^2}{(x - 3)}.$$

The points A and B are the stationary points of C .

The line $x = 3$ is a vertical asymptote to C .

a Using calculus, find the coordinates of A and B .

b Sketch the curve C^* , with equation $y = f(-x) + 2$, showing the coordinates of the images of A and B .

c State the equation of the vertical asymptote to C^* .

8

a On the same set of axes, in the interval $-\pi < \theta < \pi$, sketch the graphs of

i $y = \cot \theta$, **ii** $y = 3 \sin 2\theta$.


b Solve, in the interval $-\pi < \theta < \pi$, the equation

$$\cot \theta = 3 \sin 2\theta.$$

giving your answers, in radians, to

3 significant figures where appropriate.

9



The diagram shows, in the interval

$-\pi \leq \theta \leq \pi$, the graph of $y = k \sec(\theta - \alpha)$.

The curve crosses the y -axis at the point $(0, 4)$, and the θ -coordinate of its minimum point is $\frac{\pi}{3}$.

- a** State, as a multiple of π , the value of α .
- b** Find the value of k .
- c** Find the exact values of θ at the points where the graph crosses the line $y = -2\sqrt{2}$.
- d** Show that the gradient at the point on the curve with θ -coordinate $\frac{7\pi}{12}$ is $2\sqrt{2}$.

- 10 a** Given that $\sin^2 \theta + \cos^2 \theta \equiv 1$, show that $1 + \tan^2 \theta \equiv \sec^2 \theta$.
- b** Solve, for $0 \leq \theta < 360^\circ$, the equation $2 \tan^2 \theta + \sec \theta = 1$, giving your answers to 1 decimal place.

E

- 11 a** Prove that $\sec^4 \theta - \tan^4 \theta = 1 + 2 \tan^2 \theta$.
- b** Find all the values of x , in the interval $0 \leq x \leq 360^\circ$, for which $\sec^4 2x = \tan 2x(3 + \tan^3 2x)$. Give your answers correct to 1 decimal place, where appropriate.

- 12 a** Prove that $\cot \theta - \tan \theta = 2 \cot 2\theta$, $\theta \neq \frac{n\pi}{2}$.
- b** Solve, for $-\pi < \theta < \pi$, the equation $\cot \theta - \tan \theta = 5$, giving your answers to 3 significant figures.

- 13 a** Solve, in the interval $0 \leq \theta \leq 2\pi$ $\sec \theta + 2 = \cos \theta + \tan \theta(3 + \sin \theta)$, giving your answers to 3 significant figures.
- b** Solve, in the interval $0 \leq x \leq 360^\circ$, $\cot^2 x = \operatorname{cosec} x(2 - \operatorname{cosec} x)$, giving your answers to 1 decimal place.

- 14** Given that $y = \arcsin x$, $-1 \leq x \leq 1$ and $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$,

a express $\arccos x$ in terms of y .

b Hence find, in terms of π the value of $\arcsin x + \arccos x$.

Given that

$$y = \arccos x, \quad -1 \leq x \leq 1 \quad \text{and} \quad 0 \leq y \leq \pi,$$

- c** sketch, on the same set of axes, the graphs of $y = \arcsin x$ and $y = \arccos x$, making it clear which is which.
- d** Explain how your sketches can be used to evaluate $\arcsin x + \arccos x$.

- 15 a** By writing $\cos 3\theta$ as $\cos(2\theta + \theta)$, show that

$$\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta.$$

- b** Given that $\cos \theta = \frac{\sqrt{2}}{3}$, find the exact value of $\sec 3\theta$.

- 16** Given that $\sin(x + 30^\circ) = 2 \sin(x - 60^\circ)$,

a show that $\tan x = 8 + 5\sqrt{3}$.

b Hence express $\tan(x + 60^\circ)$ in the form $a + b\sqrt{3}$.

- 17 a** Given that $\cos A = \frac{3}{4}$ where $270^\circ < A < 360^\circ$, find the exact value of $\sin 2A$.

b i Show that

$$\cos\left(2x + \frac{\pi}{3}\right) + \cos\left(2x - \frac{\pi}{3}\right) = \cos 2x$$

Given that

$$y = 3 \sin^2 x + \cos\left(2x + \frac{\pi}{3}\right) + \cos\left(2x - \frac{\pi}{3}\right),$$

ii show that $\frac{dy}{dx} = \sin 2x$

E

- 18** Solve, in the interval $-180^\circ \leq x < 180^\circ$, the equations

a $\cos 2x + \sin x = 1$

b $\sin x(\cos x + \operatorname{cosec} x) = 2 \cos^2 x$, giving your answers to 1 decimal place.

- 19 a** Prove that

$$\frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} = 2 \operatorname{cosec} 2\theta, \quad \theta \neq 90n^\circ.$$

b Sketch the graph of $y = 2 \operatorname{cosec} 2\theta$ for $0^\circ < \theta < 360^\circ$.

c Solve, for $0^\circ < \theta < 360^\circ$, the equation

$$\frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} = 3,$$

giving your answers to 1 decimal place.

E

20 a Express $3 \sin x + 2 \cos x$ in the form $R \sin(x + \alpha)$, where $R > 0$ and $0 < \alpha < \frac{\pi}{2}$.

b Hence find the greatest value of $(3 \sin x + 2 \cos x)^4$.

c Solve, for $0 < x < 2\pi$, the equation

$$3 \sin x + 2 \cos x = 1,$$

giving your answers to 3 decimal places.

E

21 The point P lies on the curve with equation $y = \ln\left(\frac{1}{3}x\right)$.

The x -coordinate of P is 3.

Find an equation of the normal to the curve at the point P in the form $y = ax + b$, where a and b are constants.

E

22 a Differentiate with respect to x

i $3 \sin^2 x + \sec 2x$,

ii $\{x + \ln(2x)\}^3$.

Given that $y = \frac{5x^2 - 10x + 9}{(x - 1)^2}$, $x \neq -1$,

b show that $\frac{dy}{dx} = -\frac{8}{(x - 1)^3}$

E

23 Given that $y = \ln(1 + e^x)$,

a show that when $x = -\ln 3$, $\frac{dy}{dx} = \frac{1}{4}$

b find the exact value of x for which

$$e^y \frac{dy}{dx} = 6.$$

24 a Differentiate with respect to x

i $x^2 e^{3x+2}$,

ii $\frac{\cos(2x^3)}{3x}$.

b Given that $x = 4 \sin(2y + 6)$, find $\frac{dy}{dx}$ in terms of x .

E

25 Given that $x = y^2 e^{\sqrt{y}}$,

a find, in terms of y , $\frac{dx}{dy}$

b show that when $y = 4$, $\frac{dy}{dx} = \frac{e^{-2}}{12}$.

26 a Given that $y = \sqrt{1 + x^2}$, show that

$$\frac{dy}{dx} = \frac{\sqrt{3}}{2} \text{ when } x = \sqrt{3}.$$

b Given that $y = \ln\{x + \sqrt{1 + x^2}\}$, show that $\frac{dy}{dx} = \frac{1}{\sqrt{1 + x^2}}$.

27 Given that $f(x) = x^2 e^{-x}$,

a find $f'(x)$, using the product rule for differentiation


b show that $f''(x) = (x^2 - 4x + 2)e^{-x}$.

A curve C has equation $y = f(x)$.

c Find the coordinates of the turning points of C .

d Determine the nature of each turning point of the curve C .

28 a Express $(\sin 2x + \sqrt{3} \cos 2x)$ in the form $R \sin(2x + k\pi)$, where $R > 0$ and $0 < k < \frac{1}{2}$.



The diagram shows part of the curve with equation

$$y = e^{-2\sqrt{2}x}(\sin 2x + \sqrt{3} \cos 2x).$$

b Show that the x -coordinates of the turning points of the curve satisfy the equation

$$\tan\left(2x + \frac{\pi}{3}\right) = \frac{1}{\sqrt{2}}.$$

- 29** The curve C has equation $y = x^2\sqrt{\cos x}$. The point P on C has x -coordinate $\frac{\pi}{3}$.
- a** Show that the y -coordinate of P is $\frac{\sqrt{2}\pi^2}{18}$.

b Show that the gradient of C at P is 0.809, to 3 significant figures.

In the interval $0 < x < \frac{\pi}{2}$, C has a maximum at the point A .

c Show that the x -coordinate, k , of A satisfies the equation $x \tan x = 4$.


The iterative formula

$$x_{n+1} = \tan^{-1}\left(\frac{4}{x_n}\right), \quad x_0 = 1.25,$$

is used to find an approximation for k .

d Find the value of k , correct to 4 decimal places.

30



The figure shows part of the curve with equation

$$y = (2x - 1) \tan 2x, \quad 0 \leq x < \frac{\pi}{4}.$$

The curve has a minimum at the point P . The x -coordinate of P is k .

a Show that k satisfies the equation

$$4k + \sin 4k - 2 = 0.$$

The iterative formula

$$x_{n+1} = \frac{1}{4}(2 - \sin 4x_n), \quad x_0 = 0.3,$$

is used to find an approximate value for k .

- b** Calculate the values of x_1 , x_2 , x_3 and x_4 , giving your answers to 4 decimal places.
- c** Show that $k = 0.277$, correct to 3 significant figures.

E